

Diabetes Education Program

How to Fix a Low Blood Sugar

1. Check your blood sugar if:

a. Shaky, sweaty, hungry, cranky, angry, fast heartbeat, headache

2. Blood sugar less than 70?

- a. Eat a fast acting carbohydrate
 - i. Glucose tablets 3-4 **Preferred Method**
 - ii. ½ cup of juice
 - iii. ½ can regular soda
 - iv. 8-10 lifesavers
 - v. Raisins 2 tablespoons
 - vi. 8 oz milk (skim-1% is best)
- b. No high fat foods (sorry, no candy bars or doughnuts)
- c. Try not to over-treat low blood sugar
- d. Re-test your blood sugar in 15 minutes
- e. If blood sugar less than 70 repeat step 2a
- f. If blood sugar greater than 70
 - i. Eat your next planned meal
 - ii. Have a snack with 1 carbohydrate and 1 protein
 - iii. Examples:
 - 1. ½ turkey sandwich, apple with cheese, Greek yogurt with fruit

3. Talk with your doctor about low blood sugars

- a. Keep good records to determine cause
- **4.** Always keep low blood sugar supplies with you (car, purse, work, home, etc.)

5. Glucagon Emergency Kit

- a. Needed for most people on insulin (especially type 1)
- b. Family members and/or co-workers will need to learn how to give
- c. After glucagon emergency kit is given by another person:
 - i. Lay on side, with head and neck straight with spine
 - ii. Vomiting may occur
 - iii. Call 911